

## Tiavet Vet

Tiamulin Hydrogen Fumarate 450 mg/gm

## Presentation

Tiavet Vet powder: Each gram powder contains Tiamulin Hydrogen Fumarate USP 450 mg.

## Descriptions

Tiamulin is a broad spectrum antibiotic. It is specially used for prevention and treatment of Mycoplasmosis in poultry. Tiamulin inhibits bacterial protein synthesis by preventing chain formation of key enzymes and proteins.

## Indications

For the treatment & prevention of Mycoplasmosis and other infections of poultry like CRD, CCRD, Sinusitis, Airsacculitis, Enteritis, Pneumonia etc.

## Dosage & Administration

For treatment: 25-30 mg/kg body weight or 1 gm/2 Liter water for 3-5 days or 30-40 gm/100 kg feed for 5-7 days.

For prevention: 12.5-15 mg/kg body weight or 1 gm/4 Liter water for 3-7 days or 15-20 gm/100 kg feed for 3-5 days.

Or as directed by the registered Veterinarian.

## Side Effects

Tiamulin is well tolerated for poultry at recommended dosage. However, reduction of water intake may be reported.

## Precautions

Poultry should not be provided with feed containing Narasin, Monensin & Salinomycin during or for at least seven days before or after treatment with Tiamulin.

## Contraindications

Contraindicated to hypersensitivity to its any active ingredients.

#### Drug Interaction

Tiamulin should not be used with anticoccidials like Narasin, Monensin and Salinomycin.

#### Overdose

Overdose causes reduced growth and impaired feed conversion efficiency.

#### Withdrawal period

Meat : 3 days.

Egg : Not required.

#### Storage

Protect from light & keep in a cool and dry place. Keep out of the reach of children.

#### Commercial Packaging

Tiavet Vet 20 gm powder: Each box contains 1 sachet of 20 gm powder.

Tiavet Vet 100 gm powder: Each box contains 1 sachet of 100 gm powder.