

Mebastin[®]

Mebhydrolin 50 mg

Presentation

Mebastin : Each tablet contains Mebhydrolin Napadisylate INN 76 mg equivalent to Mebhydrolin 50 mg.

Description

Mebhydrolin diminishes or abolishes the main actions of histamine in the body by blocking of histamine receptors.

Indications and uses

Allergic disease or symptoms such as urticaria, pruritus of various origin, eczema, itching, drug rash, insect bites, allergic conjunctivitis, dermatitis of nutritional origin, hay fever, vasomotor rhinitis, allergic asthma.

Dosage and Administration

Adults and children over 10 years: 2-6 tablets daily

Children 5-10 years: 2-4 tablets daily

Children 2-5 years: 1-3 tablets daily

Children upto 2 years: 1-2 tablets daily

Treatment should be given in several single doses daily. Mebhydrolin may be swallowed during or shortly after meals. For children, the tablet may be crushed and mixed with food.

Contraindications

Hypersensitivity to any of the ingredients of Mebhydrolin tablet or to mebhydrolin

Prostatic hypertrophy, narrow-angle glaucoma

First trimester of pregnancy

Precautions

The substance may cause drowsiness, if affected, driving or operating machinery & alcoholic drinks should be avoided.

Side-effects

Lassitude, dizziness, hypotension, muscular weakness, nausea, vomiting, diarrhoea or constipation, epigastric pain, headache, blurred vision, tinnitus, depression, nightmares, anorexia, dryness of the mouth, tightness of the chest, tingling, sedation, drowsiness etc. may occur.

Use in pregnancy and lactation

Safety in pregnancy and lactation has not been established.

Drug interactions

The effects of atropine and tricyclic anti-depressants may be enhanced by Mebhydrolin. It may mask the warning symptoms of damage caused by ototoxic drugs.

Commercial Pack

Mebastin : Each box contains 10 blister strips of 10 tablets.

Manufactured by



Incepta Pharmaceuticals Ltd

Dhaka, Bangladesh

® Registered Trademark

MBT