

Retapex™ ointment

Retapamulin 1%

Presentation

Retapex™ ointment: Each gram ointment contains Retapamulin INN 10 mg.

Description

Retapamulin is a semi-synthetic pleuromutilin antibiotic. It is active against *Staphylococcus aureus* as well as *Streptococcus pyogenes* organisms. Retapamulin is bacteriostatic at minimum inhibitory concentration (MIC) against these pathogens. The bactericidal concentration against relevant pathogens is 1,000 times of the minimum inhibitory concentration (MIC).

Indications and Usage

Retapamulin ointment is indicated for the topical treatment of impetigo due to *Staphylococcus aureus* (methicillin-susceptible isolates only) or *Streptococcus pyogenes* in patients aged 9 months or older. It is also indicated to treat infected small lacerations, abrasions or sutured wounds.

Dosage and Administration

A thin layer of Retapamulin ointment should be applied to the affected area twice daily for 5 days. The treated area may be covered with a sterile bandage or gauze dressing if desired.

Side Effects

The most common drug related adverse reactions are application site irritation, pruritis, eczema, erythema, nasopharyngitis, headache, pyrexia, nausea, diarrhea etc.

Contraindications

None

Precautions

In the event of sensitization or severe local irritation from Retapamulin, usage should be discontinued and appropriate alternative therapy for the infection instituted. Retapamulin ointment is not intended for ingestion or for oral, intranasal, ophthalmic, or intravaginal use. Use of Retapamulin in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Pregnancy and Lactation

Pregnancy: Pregnancy Category B. It is not known whether Retapamulin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Retapamulin is administered to a nursing woman.

Lactation: The safe use of Retapamulin during breast-feeding has not been established.

Storage

Keep out of the reach of children. Store in a cool (below 30 °C) and dry place.

Commercial Pack

Retapex™ ointment: Each tube contains 10 gm of Retapamulin INN ointment.

Retapex™ ointment: Each tube contains 15 gm of Retapamulin INN ointment.



Manufactured by

Incepta Pharmaceuticals Ltd

Savar, Dhaka, Bangladesh

V.N.02

RT2