

Vinsetine®

Vinpocetine 5 mg Tablet



Presentation

Vinsetine® 5 mg Tablet: Each tablet contains Vinpocetine BP 5 mg.

Description

Vinsetine improves cerebral metabolism. It increases the glucose and oxygen consumption and utilization of the brain tissue, improves the hypoxia tolerance of the brain cell, shifts glucose metabolism towards the energetically more favorable aerobic pathway, exerts antioxidant activity, as a result of all the effects it is a cerebroprotective compound. Vinsetine considerably improves cerebral microcirculation.

Indications and uses

All forms of acute and chronic cerebral circulatory insufficiency: TIA (Transient Ischaemic Attack), reversible ischaemic neurological deficiency, progressive stroke, completed stroke, post-apoplectic conditions, multi-infarct dementia, cerebral arteriosclerosis, post traumatic, hypertensive encephalopathy etc. For the reduction of psychic or neurological symptoms of cerebral insufficiency (e.g., memory disturbance, dizziness, headache, aphasia, locomotor disorder etc.).

Ophthalmology: It can be used for the treatment of vascular disorder of the choroid and retina due to arteriosclerosis, vasospasm, macular degeneration, arterial and venous thrombosis or embolism and glaucoma secondary to above mentioned disorders.

Otology: For the treatment of impaired hearing of vascular or toxic(latrogenic) origin, presbycusis, menier's disease, cochleovestibular neuritis, tinnitus and dizziness of labyrinth origin.

Dosage guideline

Vinsetine is taken 1-2 tablets three times daily with meals. The maintenance dose is 1 tablet three times daily over long periods.

Side-effects

Vinsetine is well tolerated. In some cases transient fall of blood pressure and tachycardia may occur.

Contraindications

Vinpocetine is contraindicated in pregnancy and lactation. Hypersensitivity to its components is also contraindicated.

Precaution

Because Vinsetine can reduce the ability of blood to clot, those individuals with a tendency to bleed should avoid Vinpocetine.

Drug interaction

Should not be given with Heparin.

Storage condition

Do not store above 30 °C. Keep away from light and out of the reach of children.

Commercial Pack

Vinsetine® 5 mg Tablet: Each box contains 10 blister strips of 10 tablets.

Manufactured by
 **Incepta Pharmaceuticals Ltd**
Savar, Dhaka, Bangladesh
® Registered Trademark



V.N.01

VNT