

Reset[®]

Paracetamol Tablet, Syrup & Paediatric drops

Presentation

Reset[®] : Each tablet contains Paracetamol BP 500 mg.

Reset[®] Syrup: Each 5 ml syrup contains Paracetamol BP 120 mg.

Reset[®] Paediatric Drops: Each ml paediatric drops contains Paracetamol BP 80 mg.

Description

Reset (Paracetamol) is a fast acting and safe analgesic with marked antipyretic property. It is specially suitable for patients who, for any reason, can not tolerate aspirin or other analgesics.

Indications

All conditions requiring relief from pain and fever such as neuritis, neuralgia, headache, toothache, earache, bodyache, myalgia, minor arthritis pain, pain due to rheumatic disorder, cold, influenza, dysmenorrhoea etc..

Dosage and Administration

Tablet

Adult: 1 - 2 tablets every 4 to 6 hours up to a maximum of 4 g (8 tablets) daily

Children (6 - 12 years): ½ to 1 tablet 3 to 4 times daily

Syrup

Children:

- 2 months: 60 mg (½ measuring spoonful) for post immunization pyrexia
- Under 3 months (on doctor's advice only): 10 mg/kg (5 mg/kg if jaundiced)
- 3 months - 1 year: 60 - 120 mg (½ - 1 measuring spoonful)
- 1 - 5 years: 1 - 2 measuring spoonful
- 6 - 12 years: 2 - 4 measuring spoonful

Paediatric drops

2 months: 60 mg for post-immunization pyrexia; under 3 months (on doctor's advice only), 10 mg/kg (5 mg/kg if jaundiced); 3 months - 1 year: 60-120 mg; 1 - 5 years: 120-250 mg.

Paediatric drops may be given to children and neonates according to following guideline:

Age	Dose of Paediatric Drops	Frequency
0-3 months	0.5 ml	Upto 4 times daily
4-11 months	1 ml	
12-23 months	1.5 ml	
2-3 years	2 ml	
4-5 years	3 ml	

Side-effects

Side effects of paracetamol are usually mild, though haematological reactions including thrombocytopenia, leukopenia, pancytopenia, neutropenia, and agranulocytosis have been reported. Pancreatitis, skin rashes, and other allergic reactions occur occasionally.

Precautions

Paracetamol should be given with care to patients with impaired kidney or liver function. Paracetamol should be given with care to patients taking other drugs that affect the liver.

Contraindications

Paracetamol is contraindicated in patients with severe renal function impairment, hepatic disease (viral hepatitis) and known hypersensitivity to paracetamol.

Overdosage

Symptoms of Paracetamol overdose in the first 24 hours are pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 40 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur.

Storage

Do not store above 30°C. Keep away from light and out of the reach of children.

Commercial Pack


Reset[®] : Box containing 30 blister strips of 10 tablets.

Reset[®] Syrup: Bottle containing 60 ml syrup.

Reset[®] Syrup: Bottle containing 100 ml syrup.

Reset[®] Paediatric Drops: Bottle containing 15 ml Paediatric Drops.

Manufactured by

 **Incepta Pharmaceuticals Ltd**

Savar, Dhaka, Bangladesh

® Registered Trademark

V.N.02

RSD